

New QCD Sum Rules for Nucleon Tensor Charge*

Xuemin Jin[†] and Jian Tang[‡]

Center for Theoretical Physics

Laboratory for Nuclear Science

and Department of Physics

Massachusetts Institute of Technology

Cambridge, Massachusetts 02139, USA

(MIT-CTP-2634 hep-th/9705269 May 1997)

Abstract

Two new QCD sum rules for nucleon tensor charge are derived from a mixed correlator of spin-1/2 and spin-3/2 nucleon interpolating fields. These sum rules are analyzed along with a sum rule obtained from the usual correlator of a general spin-1/2 nucleon interpolating field. The validity and reliability of the sum rules are examined by monitoring the contaminations arising from the transitions and continuum and the convergence of the operator product expansion. Valid sum rules are identified and their predictions are presented. It is found that the vacuum tensor susceptibility induced by the external field plays an important role in determining both the validity and predictions of the sum rules. The uncertainties associated with the sum-rule predictions are also discussed.

Submitted to: *Physical Review D*

[†]jin@ctpa02.mit.edu

[‡]jtang@ctpa02.mit.edu

*This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement #DF-FC02-94ER40818.

I. INTRODUCTION

Tensor charge of the nucleon, defined through the nucleon matrix element of the tensor current, $\bar{q}\sigma_{\alpha\beta}q$, at zero momentum transfer, is perhaps the least known nucleon charge both theoretically and experimentally. Like other charges of the nucleon, the tensor charge reveals important information on the nonperturbative structure of the nucleon and understanding it from the underlying theory of strong interaction, quantum chromodynamics (QCD), is of great importance.

The nucleon tensor charge δq is defined by

$$\langle N | \bar{q} \sigma_{\alpha\beta} q | N \rangle = \delta q \bar{U} \sigma_{\alpha\beta} U , \quad (1)$$

where U stands for the nucleon Dirac spinor. It has been shown by Jaffe and Ji [1] that the first moment of the twist-two transversity quark distribution in a nucleon, $h_1(x)$, is related to the nucleon tensor charge

$$\int_0^1 dx \left[h_1^q(x) - \bar{h}_1^q(x) \right] = \delta q , \quad (2)$$

where h_1^q and $\bar{h}_1^q$ receive contributions from quark and antiquark (of flavor q), respectively. Unlike the twist-two spin averaged quark distribution $f_1(x)$ and helicity difference quark distribution $g_1(x)$, the $h_1(x)$ flips chirality and is thus suppressed in inclusive deep inelastic lepton scattering [2]. As such, there is no existing experimental data on $h_1(x)$ and the nucleon tensor charge. However, there is no such suppression of $h_1(x)$ in deep inelastic processes with hadronic initial states such as Drell-Yan [3,2], and experiments have been planned to measure the quark transversity distribution in the nucleon at RHIC [4], HERA [5], and CERN [6].

Recently, the nucleon tensor charge has been investigated theoretically in various models and approaches [7–11]. Previous studies of the nucleon tensor charge via the QCD sum-rule method have been made by He and Ji [7,8]. These authors have found that the sum rules obtained from the correlator of Ioffe’s nucleon interpolating field [12] are unstable, making the extraction of the nucleon tensor charge from those sum rules difficult. In this paper, we derive two new QCD sum rules from the mixed correlator of spin-1/2 and spin-3/2 interpolating fields using the external-field method. The consideration of the mixed correlator is motivated by its success in nucleon mass sum rules [13] and in the sum rules for nucleon axial-vector coupling constants [14]. We analyze these new sum rules along with a sum rule obtained from the usual correlator with a general spin-1/2 nucleon interpolating field. We examine the performances and reveal the validity and reliability of these sum rules.

A well-known problem that arises in calculating hadronic matrix elements within the QCD sum-rule method is the unwanted physics associated with transitions from ground state to excited states. The contributions of these transitions are not exponentially suppressed relative to the ground state contribution which contains the ground state property of interest [15]. In practice, there are two formalisms for treating the transitions. The usual formalism is to model the sum over all the transitions from ground state to excited states by a single unknown parameter. This parameter is extracted from the sum rules, along with the ground state property, and the continuum threshold is usually fixed at the same value as that for

the corresponding mass sum rule. The resolution of the ground state signal relies on the different polynomial (in Borel mass) behaviors of the ground state signal and the transition term.

An alternative formalism for treating the transitions is to multiply the invariant functions by the factor $M_h^2 - q^2$ (with M_h the ground state hadron mass) before performing the Borel transform. The transition contribution is then exponentially suppressed relative to the ground state contribution and can be included in the continuum model in practice. In this case, it is important to treat the continuum threshold as an unknown parameter to be extracted from the sum rule along with the ground state property. Assuming the threshold to be the same as that for the mass sum rule is unjustified and may introduce artificial bias to the extracted hadron properties [16]. We observe that in the previous works [7,8] the alternative formalism was adopted but the continuum threshold was taken to be the same as the one used in the nucleon mass sum rule. In the present paper, we shall consider both formalisms.

Both sides of QCD sum rules depend on an auxiliary parameter—Borel mass, which is introduced through the use of the Borel transform. If a sum rule were perfect, one would expect that the two sides of the sum rule overlap for all values of the Borel mass. However, in practice, the two sides of the sum rules overlap only in a limited range of the Borel mass (at best) because of the truncation of the operator product expansion (OPE) and the crudity of the models for the transitions and continuum. Thus, in order to extract the properties of the ground state by matching the sum rules, one should work in a region of Borel mass where the ground state contribution dominates the phenomenological side. This usually sets an upper bound in the Borel mass space, beyond which the excited-state and transition contributions dominate and the background noise is hence too strong to reliably isolate the ground state signal. On the other hand, the truncated OPE must be sufficiently convergent as to accurately describe the true OPE. This, in practice, sets a lower limit in the Borel mass space, beyond which higher order terms not present in the truncated OPE may be out of control.

Therefore, we expect a sum rule to work if the two sides of the sum rule match in a *valid* window in Borel mass space where the ground state contribution of interest dominates the phenomenological side and the higher order OPE terms are under control. The validity and reliability of a sum rule is then determined by the quality of overlap of the two sides, the size of valid Borel window, and the relative contributions of the excited states and the highest OPE terms in the valid Borel window. A sum rule failing to have a valid Borel window should be considered invalid, and results obtained from such an invalid sum rule may be meaningless and misleading. In this paper, we will analyze the sum rules with respect to the above criteria.

This paper is organized as follows. In Sec. II, we derive QCD sum rules for the nucleon tensor charge within the external-field approach. In Sec. III, we analyze the sum rules and present results and discussions. Section IV is devoted to summary and conclusion.

II. SUM RULES FOR NUCLEON TENSOR CHARGE

In this section, we first give a brief outline of the external-field QCD sum-rule method with an emphasis on some issues related to the treatment of the transitions between ground state and excited states. We then use this method to derive QCD sum rules for the nucleon tensor charge.

A. Outline of the method

Consider the nucleon two-point correlator in the presence of a constant external (anti-symmetric) tensor field, $Z^{\alpha\beta}$, defined by

$$\Pi(q) = i \int d^4x e^{iq \cdot x} \langle 0 | T \eta(x) \bar{\eta}(0) | 0 \rangle_Z , \quad (3)$$

where η is a nucleon interpolating field, constructed from quark fields, which couples to the nucleon. To lowest dimension, there are two independent interpolating fields which couple to spin-1/2 states only and carry the quantum numbers of the nucleon. So, a general spin-1/2 interpolating field for the proton can be expressed as

$$\eta^{1/2} = \eta_1 + \beta \eta_2 , \quad (4)$$

where β is a real parameter and

$$\eta_1(x) = \epsilon_{abc} \left[u^{aT}(x) C \gamma^5 d^b(x) \right] u^c(x) , \quad (5)$$

$$\eta_2(x) = \epsilon_{abc} \left[u^{aT}(x) C d^b(x) \right] \gamma_5 u^c(x) . \quad (6)$$

Here $u(x)$ and $d(x)$ denote up and down quark operators, a, b , and c are color indices, and C is the charge conjugation matrix. The interpolating field for the neutron can be obtained by changing $u(d)$ into $d(u)$. The interpolating field advocated by Ioffe [12] and used in previous sum-rule calculations of the nucleon tensor charge [7,8] may be recovered by setting $\beta = -1$ and multiplying an overall factor of -2 .

There is a spin-3/2 interpolating field which also couples to the proton through its spin-1/2 component [17,13]

$$\eta_\mu^{3/2} = \epsilon_{abc} \left[\left(u^{aT} C \sigma_{\rho\lambda} d^b \right) \sigma^{\rho\lambda} \gamma_\mu u^c - \left(u^{aT} C \sigma_{\rho\lambda} u^b \right) \sigma^{\rho\lambda} \gamma_\mu d^c \right] . \quad (7)$$

The couplings of $\eta^{1/2}$ and $\eta_\mu^{3/2}$ to the proton are defined as

$$\langle 0 | \eta^{1/2} | N \rangle = \lambda_1 U(q) , \quad \langle N | \bar{\eta}_\mu^{3/2} | 0 \rangle = \lambda_2 \bar{U}(q) \gamma_5 \left(\frac{4q_\mu}{M_N} + \gamma_\mu \right) , \quad (8)$$

where $U(q)$ denotes the Dirac spinor of the proton with the normalization $\bar{U}(q)U(q) = 2M_N$, and λ_1 and λ_2 describe the coupling strengths of the interpolating fields to the proton.

Therefore, independent QCD sum rules can be obtained from the correlators of various combinations of $\eta^{1/2}$ and $\eta_\mu^{3/2}$. In the present work, we will consider the mixed correlator

of $\eta^{1/2}$ and $\eta_\mu^{3/2}$ as well as the $1/2$ correlator of two $\eta^{1/2}$'s. While the latter, at $\beta = -1$, leads to the sum rules discussed previously, the former gives rise to new independent sum rules for the nucleon tensor charge. In Ref [13], it has been demonstrated that the nucleon mass sum rules from the mixed correlator have broader valid Borel windows and relatively smaller continuum contamination as compared to the sum rules from the $1/2$ correlator.

The subscript Z in Eq. (3) denotes that $\Pi(q)$ is evaluated by adding a tensor coupling term

$$\Delta\mathcal{L} = Z^{\alpha\beta} \sum_q g_q \bar{q}(x) \sigma_{\alpha\beta} q(x) , \quad (9)$$

to the QCD Lagrangian, where $q = \{u, d, s\}$. Here g_q keeps track of the flavor structure of the external field. For example, if one is interested in the isovector (isoscalar) tensor charge, $g_\tau^v \equiv \delta u - \delta d$ ($g_\tau^s \equiv \delta u + \delta d$), one simply sets $g_u = -g_d = 1$ ($g_u = g_d = 1$) and $g_s = 0$. The external-field method proceeds by expanding $\Pi(q)$ to first order in the external field

$$\Pi(q) = \Pi^{(0)}(q) + Z^{\alpha\beta} \Pi_{\alpha\beta}^{(1)}(q) + \cdots , \quad (10)$$

where $\Pi^{(0)}(q)$ is the correlator in the absence of the external field from which the usual nucleon mass sum rules are derived. It is the linear response to the external field, $\Pi_{\alpha\beta}^{(1)}(q)$, that gives rise to the sum rules for the nucleon tensor charge.

The QCD side of the sum rule is obtained by carrying out the OPE. The external field can interact with the quark field directly; it also polarizes the QCD vacuum and induces nonperturbative condensates. The latter are summarized by vacuum susceptibilities, which, to first order in the external field, are defined as [7,8]

$$\langle \bar{q} \sigma_{\alpha\beta} q \rangle_Z \equiv g_q \chi \langle \bar{q} q \rangle Z_{\alpha\beta} , \quad (11)$$

$$\langle \bar{q} g_s G_{\alpha\beta} q \rangle_Z \equiv g_q \kappa \langle \bar{q} q \rangle Z_{\alpha\beta} , \quad (12)$$

$$\langle \bar{q} g_s \tilde{G}_{\alpha\beta} \gamma_5 q \rangle_Z \equiv -i g_q \xi \langle \bar{q} q \rangle Z_{\alpha\beta} , \quad (13)$$

where g_s is the strong coupling constant ($D_\mu = \partial_\mu + i g_s \mathcal{A}_\mu$), $\tilde{G}_{\alpha\beta} \equiv \frac{1}{2} \epsilon_{\alpha\beta\rho\lambda} G^{\rho\lambda}$ with $G_{\alpha\beta}$ the gluon field tensor, χ , κ , and ξ are vacuum susceptibilities, and $\langle \hat{O} \rangle$ denotes the usual vacuum condensate. Here, the isospin breaking effect has been neglected and will be henceforth. The strangeness contribution will also be ignored in the present paper as it is expected to be small [9,11]. The calculation of the Wilson coefficients is straightforward following the techniques discussed extensively in the literature. Some details of the calculations are presented in the Appendix.

We now turn to the phenomenological description, which can be obtained by expanding $\Pi_{\alpha\beta}^{(1)}(q)$ in terms of physical intermediate states. For a generic invariant function, the phenomenological representation can be written as

$$\begin{aligned} \Pi^{(1)}(q^2) \sim & \frac{\lambda_{\mathcal{O}}^2 g_\tau}{(M_N^2 - q^2)^2} + \sum_{N^*} \frac{C_{N \leftrightarrow N^*}}{(M_N^2 - q^2)(M_{N^*}^2 - q^2)} \\ & + \text{terms involving only excited states} , \end{aligned} \quad (14)$$

where M_N and M_{N^*} are the masses of the ground state nucleon and the excited state, respectively, and $\lambda_{\mathcal{O}}^2 = \lambda_1 \lambda_2$, λ_1^2 for the mixed and $1/2$ correlators, respectively. The first term contains the nucleon tensor charge $g_T = \sum_q g_q \delta q$ of interest, the second term is the transition term from ground state to excited states, and the rest involves only the excited states. After a usual Borel transformation, one finds

$$\Pi^{(1)}(M^2) \sim \frac{\lambda_{\mathcal{O}}^2}{M^2} g_T e^{-M_N^2/M^2} + \sum_{N^*} \left[\frac{C_{N \leftrightarrow N^*}}{M_{N^*}^2 - M_N^2} \left(1 - e^{-(M_{N^*}^2 - M_N^2)/M^2} \right) \right] e^{-M_N^2/M^2} \\ + \text{exponentially suppressed terms} , \quad (15)$$

where M is the Borel mass. It can be seen clearly that the second term is not exponentially damped as compared to the first term. The usual formalism is to approximate the second term as $A e^{-M_N^2/M^2}$ with A a phenomenological constant to be extracted from the sum rule along with g_T .

An alternative formalism is to use the combination $(M_N^2 - q^2) \Pi^{(1)}(q^2)$, instead of $\Pi^{(1)}(q^2)$ alone. The phenomenological side then becomes

$$\tilde{\Pi}^{(1)}(M^2) \sim \lambda_{\mathcal{O}}^2 g_T e^{-M_N^2/M^2} + \sum_{N^*} C_{N \leftrightarrow N^*} e^{-M_{N^*}^2/M^2} + \text{exponentially suppressed terms} . \quad (16)$$

The transitions from ground state to excited states (second term) are now exponentially suppressed with respect to the ground state (first) term and hence can be absorbed into the continuum model. As such, one no longer needs to introduce any phenomenological parameter to represent the transitions from the ground state to excited states as their contribution has been included in the continuum model. However, as stressed in Ref. [16] the continuum threshold must be treated as an unknown parameter to be extracted from the sum rule.

Since the linear response $\Pi_{\alpha\beta}^{(1)}$ in general contains distinct Dirac and Lorentz structures, one may obtain many sum rules, one for each invariant structure. However, these sum rules do not work equally well in practice. In particular, some sum rules work well while the others may fail. Recently, it has been pointed out by the present authors [18] that chirality plays an important role in determining the reliability of baryon sum rules. There it was argued that for light baryons the chiral-odd sum rules (where the chiral-odd operators dominate) are generally more reliable than the chiral-even sum rules (where the chiral-even operators dominate). Thus, we will only consider the chiral-odd sum rules and disregard the chiral-even sum rules in the discussions to follow.

B. Sum rules from the mixed correlator

The mixed correlator is defined by

$$\Pi_{\mu\nu}(q) = i \int d^4x e^{iq \cdot x} \langle 0 | T [\eta_\mu^{1/2}(x) \bar{\eta}_\nu^{3/2}(0)] | 0 \rangle_Z \\ = \Pi_{\mu\nu}^{(0)}(q) + Z^{\alpha\beta} \Pi_{\mu\nu, \alpha\beta}^{(1)}(q) + \dots , \quad (17)$$

where $\eta_\mu^{1/2} \equiv \gamma_\mu \gamma_5 \eta^{1/2}$ and $\eta^{1/2}$ and $\eta_\nu^{3/2}$ are given in Eqs. (4) and (7), respectively. The use of $\eta_\mu^{1/2}$ is to facilitate the calculations and does not affect the sum rules. The linear response of $\Pi_{\mu\nu}(q)$ to the external field can be decomposed into various invariant structures:

$$\begin{aligned} \Pi_{\mu\nu,\alpha\beta}^{(1)}(q) = & \Pi_1(q^2) \gamma_\mu (\hat{q} \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \hat{q}) \gamma_\nu + \Pi_2(q^2) \gamma_\mu \sigma_{\alpha\beta} q_\nu \\ & + \Pi_3(q^2) i \gamma_\mu (q_\alpha \gamma_\beta - q_\beta \gamma_\alpha) \hat{q} q_\nu + \cdots, \end{aligned} \quad (18)$$

where $\hat{q} \equiv q_\mu \gamma^\mu$. Here, the first three structures receive contributions from the ground state proton and lead to chiral-odd sum rules. Since there is no continuum model for the sum rule from Π_3 , we only focus on the sum rules from Π_1 and Π_2 . Adopting the usual formalism for the transitions, we obtain the sum rule from Π_1

$$\begin{aligned} & \frac{a}{8} c_1 E_0 L^{4/9} M^2 - \frac{m_0^2 a}{24} c_2 L^{-2/27} \left[\ln \left(\frac{M^2}{\Lambda^2} \right) - \gamma_{\text{EM}} \right] + \frac{m_0^2 a}{96} c_3 L^{-2/27} \\ & - \frac{\chi a^2}{24} c_4 L^{8/27} - \frac{\kappa a^2}{144 M^2} c_5 L^{8/9} - \frac{\xi a^2}{72 M^2} c_6 L^{8/9} + \frac{\chi m_0^2 a^2}{1728 M^2} c_7 L^{-6/27} \\ & + \frac{ab}{864 M^2} c_8 L^{4/9} = M_N \tilde{\lambda}_1 \tilde{\lambda}_2 \left(\frac{g_{\text{T}}}{M^2} + A_1 \right) e^{-M_N^2/M^2}, \end{aligned} \quad (19)$$

and the sum rule from Π_2

$$\begin{aligned} & \frac{2a}{3} d_1 E_0 L^{4/9} M^2 - \frac{m_0^2 a}{3} d_2 L^{-2/27} \left[\ln \left(\frac{M^2}{\Lambda^2} \right) - \gamma_{\text{EM}} \right] - \frac{m_0^2 a}{6} d_3 L^{-2/27} \\ & + \frac{\chi a^2}{3} d_4 L^{8/27} + \frac{\kappa a^2}{9 M^2} d_5 L^{8/9} + \frac{\xi a^2}{9 M^2} d_6 L^{8/9} - \frac{\chi m_0^2 a^2}{216 M^2} d_7 L^{-6/27} \\ & + \frac{ab}{216 M^2} d_8 L^{4/9} = 4 \frac{\tilde{\lambda}_1 \tilde{\lambda}_2}{M_N} \left[g_{\text{T}} \left(\frac{2M_N^2}{M^2} - 1 \right) + A_2 \right] e^{-M_N^2/M^2}, \end{aligned} \quad (20)$$

where $a = -(2\pi)^2 \langle \bar{q} q \rangle$, $b = \langle g_s^2 G^2 \rangle$, $m_0^2 = -\langle \bar{q} g_s \sigma \cdot G q \rangle / \langle \bar{q} q \rangle$, $\tilde{\lambda}_1 = (2\pi)^2 \lambda_1$, $\tilde{\lambda}_2 = (2\pi)^2 \lambda_2$, $E_0 = 1 - e^{-s_0/M^2}$ with s_0 the continuum threshold, and $\gamma_{\text{EM}} = 0.577 \cdots$ the Euler-Mascheroni constant. The anomalous dimensions of various operators have been taken into account through the factor $L = \ln(M^2/\Lambda_{\text{QCD}}^2) / \ln(\mu^2/\Lambda_{\text{QCD}}^2)$, where $\mu = 500 \text{ MeV}$ is the renormalization scale and $\Lambda_{\text{QCD}} = 150 \text{ MeV}$ is the QCD scale parameter. Here Λ denotes the infrared cut-off, which arises from the quark propagator in the presence of both the external field and background gluon field (see Appendix); a reasonable choice for Λ is $\Lambda = \mu$ [19]. The c_i and d_i are defined as

$$\begin{aligned}
c_1 &= (5 + 9\beta)g_u + (1 - 9\beta)g_d, & c_2 &= (1 - \beta)(4g_u - g_d), \\
c_3 &= -(29 + 85\beta)g_u + (5 - 71\beta)g_d, & c_4 &= -7(1 + \beta)g_u + (1 - 11\beta)g_d, \\
c_5 &= 69\beta g_u + (-6 + 2\beta)g_d, & c_6 &= (7 + 9\beta)g_u - (1 + 21\beta)g_d, \\
c_7 &= -(47 + 87\beta)g_u + (23 - 237\beta)g_d, & c_8 &= 5(1 + \beta)g_u - (8 - 28\beta)g_d, \\
d_1 &= (3 + 4\beta)g_u + 5\beta g_d, & d_2 &= (1 - \beta)(g_u - g_d), \\
d_3 &= (2 + 13\beta)g_u + (1 + 11\beta)g_d, & d_4 &= (7 + 3\beta)g_u - (1 - 3\beta)g_d, \\
d_5 &= (10 - 13\beta)g_u - (4 + 2\beta)g_d, & d_6 &= 2\beta g_u + 6\beta g_d, \\
d_7 &= -(158 + 10\beta)g_u + (7 + 71\beta)g_d, & d_8 &= (5 + 7\beta)g_u + (1 + 23\beta)g_d.
\end{aligned} \tag{21}$$

When the alternative formalism for the transitions is used, the sum rule from Π_1 is

$$\begin{aligned}
& \frac{a}{8} c_1 (M_N^2 M^2 E_0 - M^4 E_1) L^{4/9} - \frac{m_0^2 a}{24} c_2 \left[M_N^2 \ln \left(\frac{M^2}{\Lambda^2} \right) - M_N^2 \gamma_{\text{EM}} - M^2 E_0 \right] L^{-2/27} \\
& + \frac{m_0^2 a}{96} c_3 M_N^2 L^{-2/27} - \frac{\chi a^2}{24} c_4 M_N^2 L^{8/27} - \frac{\kappa a^2}{144} c_5 \left(1 + \frac{M_N^2}{M^2} \right) L^{8/9} \\
& - \frac{\xi a^2}{72} c_6 \left(1 + \frac{M_N^2}{M^2} \right) L^{8/9} + \frac{\chi m_0^2 a^2}{1728} c_7 \left(1 + \frac{M_N^2}{M^2} \right) L^{-6/27} \\
& + \frac{ab}{864} c_8 \left(1 + \frac{M_N^2}{M^2} \right) L^{4/9} = M_N \tilde{\lambda}_1 \tilde{\lambda}_2 g_T e^{-M_N^2/M^2},
\end{aligned} \tag{22}$$

and the sum rule from Π_2 is

$$\begin{aligned}
& \frac{2a}{3} d_1 (M_N^2 M^2 E_0 - M^4 E_1) L^{4/9} - \frac{m_0^2 a}{3} d_2 \left[M_N^2 \ln \left(\frac{M^2}{\Lambda^2} \right) - M_N^2 \gamma_{\text{EM}} - M^2 E_0 \right] L^{-2/27} \\
& - \frac{m_0^2 a}{6} d_3 M_N^2 L^{-2/27} + \frac{\chi a^2}{3} d_4 M_N^2 L^{8/27} + \frac{\kappa a^2}{9} d_5 \left(1 + \frac{M_N^2}{M^2} \right) L^{8/9} \\
& + \frac{\xi a^2}{9} d_6 \left(1 + \frac{M_N^2}{M^2} \right) L^{8/9} - \frac{\chi m_0^2 a^2}{216} d_7 \left(1 + \frac{M_N^2}{M^2} \right) L^{-6/27} \\
& + \frac{ab}{216} d_8 \left(1 + \frac{M_N^2}{M^2} \right) L^{4/9} = 8 M_N \tilde{\lambda}_1 \tilde{\lambda}_2 g_T e^{-M_N^2/M^2},
\end{aligned} \tag{23}$$

where $E_1 = 1 - e^{-s_0/M^2} (s_0/M^2 + 1)$.

C. Sum rules from the 1/2 correlator

The 1/2 correlator is given by

$$\begin{aligned}
\Pi(q) &= i \int d^4x e^{iq \cdot x} \langle 0 | T[\eta^{1/2}(x) \bar{\eta}^{1/2}(0)] | 0 \rangle_z \\
&= \Pi^{(0)}(q) + Z^{\alpha\beta} \Pi_{\alpha\beta}^{(1)}(q) + \dots .
\end{aligned} \tag{24}$$

The linear response term in this case has three distinct structures

$$\Pi_{\alpha\beta}^{(1)} = \Pi_4(q^2) (\hat{q} \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \hat{q}) + \Pi_5(q^2) \sigma_{\alpha\beta} + \Pi_6(q^2) i (q_\alpha \gamma_\beta - q_\beta \gamma_\alpha) \hat{q} . \tag{25}$$

However, only Π_4 leads to a chiral-odd sum rule, which is given by

$$\begin{aligned}
&\frac{a}{32} f_1 L^{4/27} E_0 M^2 + \frac{m_0^2 a}{96} f_2 \left[\ln \left(\frac{M^2}{\Lambda^2} \right) - \gamma_{\text{EM}} \right] L^{-10/27} - \frac{m_0^2 a}{192} f_3 L^{-10/27} \\
&- \frac{\chi a^2}{48} f_4 + \frac{\kappa a^2}{576 M^2} f_5 L^{16/27} + \frac{\xi a^2}{96 M^2} f_6 L^{16/27} - \frac{\chi m_0^2 a^2}{1728 M^2} f_7 L^{-14/27} \\
&+ \frac{ab}{6912 M^2} f_8 L^{4/27} = \tilde{\lambda}_1^2 \left(\frac{M_N}{M^2} g_{\text{T}} + A_4 \right) e^{-M_N^2/M^2} ,
\end{aligned} \tag{26}$$

in the usual formalism and

$$\begin{aligned}
&\frac{a}{32} f_1 L^{4/27} (M_N^2 E_0 M^2 - E_1 M^4) + \frac{m_0^2 a}{96} f_2 \left[M_N^2 \ln \left(\frac{M^2}{\Lambda^2} \right) - M_N^2 \gamma_{\text{EM}} - E_0 M^2 \right] L^{-10/27} \\
&- \frac{m_0^2 a}{192} f_3 M_N^2 L^{-10/27} - \frac{\chi a^2}{48} f_4 M_N^2 + \frac{\kappa a^2}{576} f_5 L^{16/27} \left(1 + \frac{M_N^2}{M^2} \right) \\
&+ \frac{\xi a^2}{48} f_6 L^{16/27} \left(1 + \frac{M_N^2}{M^2} \right) - \frac{\chi m_0^2 a^2}{1728} f_7 L^{-14/27} \left(1 + \frac{M_N^2}{M^2} \right) \\
&+ \frac{ab}{6912} f_8 L^{4/27} \left(1 + \frac{M_N^2}{M^2} \right) = M_N \tilde{\lambda}_1^2 g_{\text{T}} e^{-M_N^2/M^2} ,
\end{aligned} \tag{27}$$

in the alternative formalism, where

$$\begin{aligned}
f_1 &= (10 + 6\beta^2) g_u - (3 + 2\beta - \beta^2) g_d , & f_2 &= (1 - \beta)^2 g_u , \\
f_3 &= (4 - 10\beta + 6\beta^2) g_u - 5(1 - \beta^2) g_d , & f_4 &= -2(1 - \beta) g_u + (1 - \beta^2) g_d , \\
f_5 &= (20 + 2\beta - 22\beta^2) g_u + 9(1 - \beta^2) g_d , & f_6 &= 2(1 - \beta) g_u - (1 - \beta^2) g_d , \\
f_7 &= (6 - 7\beta + \beta^2) g_u - 16(1 - \beta^2) g_d , \\
f_8 &= (62 + 36\beta + 70\beta^2) g_u + (1 - 6\beta + 17\beta^2) g_d .
\end{aligned} \tag{28}$$

The sum rule Eq. (27), at $\beta = -1$, can be compared with Eq. (20) of Ref. [8]. While the first and fourth terms agree, the rest disagree. The factor $M_N^2 \ln(M^2/\Lambda^2) - M_N^2 \gamma_{\text{EM}} - M^2 E_0$ in the second term has the form $M_N^2 \ln(M_N^2/\Lambda^2) - M_N^2$ in Ref. [8]. We have also compared the corresponding form of this term before the Borel transformation and found an overall sign difference. Furthermore, the Borel transformation of this term has not been done

correctly in Ref. [8] (see Appendix for more details). The third term is absent in Ref. [8]. The differences in the last four terms arise from the incomplete calculation of the Wilson coefficients in Ref. [8] (see Appendix). There is also a minor difference in the anomalous dimension for the mixed quark-gluon operator $\bar{q}g_s\sigma\cdot Gq$. We use $-2/27$, while Ref. [8] simply takes 0.

III. SUM-RULE ANALYSIS

We now analyze the sum rules derived in the previous section. To this end, we optimize the fit of the two sides of each individual sum rule in a valid Borel window, which is chosen such that the highest order (in $1/M^2$) term contributes no more than $\sim 10\%$ to the QCD side while the contribution of the transition plus continuum is less than 50% of the total phenomenological side (i.e., the sum of the contributions from ground state, transition, and continuum). Those sum rules which fail to have a valid Borel window is considered invalid and useless. Such criteria have been adopted in various sum-rule calculations [13,14,20]. While the selection of 50% is obvious for the ground state dominance, the selection of 10% is a reasonably conservative criterion. To quantify the fit of the two sides, we sample the sum rule in the valid Borel window and minimize the measure $\delta(M^2) = (\text{LHS-RHS})^2$ averaged over 50 points evenly spaced in the valid Borel window. Here, LHS and RHS denote the left- and right-hand sides of the sum rule, respectively.

A. Inputs

The inputs for the sum rules can be classified into three categories: a) the usual vacuum condensates; b) the vacuum susceptibilities χ , κ , and ξ ; c) the nucleon mass M_N , the coupling strengths λ_1 and λ_2 , and the continuum threshold s_0 . [The continuum threshold is needed as input for the sum rules (19–20) and (26).] For the values of vacuum condensates we use the central values given in Ref. [13]: $a = 0.52 \text{ GeV}^3$, $b = 1.2 \text{ GeV}^4$, and $m_0^2 = 0.72 \text{ GeV}^2$. The mixing parameter β can be chosen to minimize the overlap with the excited states and broaden the valid Borel window. Here we adopt $\beta = 0$ for the sum rules from the mixed correlator and $\beta = -1.2$ for the sum rules from the $1/2$ correlator, as determined in Ref. [13].

The vacuum susceptibilities arise from the response of condensates to the external field. It is easy to show that to first order in the external field

$$\langle \bar{q}\mathcal{O}_{\alpha\beta}q \rangle_Z = Z^{\gamma\delta} \lim_{p\lambda \rightarrow 0} i \int d^4x e^{ip\cdot y} \langle T[\sum_q g_q \bar{q}(y) \sigma_{\gamma\delta} q(y), \mathcal{O}_{\alpha\beta}(0)] \rangle, \quad (29)$$

where $\mathcal{O}_{\alpha\beta} = \{\bar{q}\sigma_{\alpha\beta}q, \bar{q}g_s G_{\alpha\beta}q, \bar{q}g_s \tilde{G}\gamma_5 q\}$. The two-point correlation function in Eq. (29) can be decomposed into

$$\begin{aligned} i \int d^4x e^{ip\cdot y} \langle T[\sum_q g_q \bar{q}(y) \sigma_{\gamma\delta} q(y), \mathcal{O}_{\alpha\beta}(0)] \rangle &= (g_{\gamma\alpha} g_{\delta\beta} - g_{\gamma\beta} g_{\delta\alpha}) \Pi_1^{\mathcal{O}}(p^2) \\ &+ (g_{\gamma\alpha} p_\delta p_\beta - g_{\delta\alpha} p_\gamma p_\beta - g_{\gamma\beta} p_\delta p_\alpha + g_{\delta\beta} p_\gamma p_\alpha) \Pi_2^{\mathcal{O}}(p^2). \end{aligned} \quad (30)$$

Neglecting isospin breaking effect and strangeness contribution and using Eqs. (11–13), we find

$$\chi a = -8\pi^2 \Pi_1^\chi(0) , \quad \kappa a = -8\pi^2 \Pi_1^\kappa(0) , \quad \xi a = -8\pi^2 i \Pi_1^\xi(0) . \quad (31)$$

These definitions are equivalent to those given in Ref. [8]. It is worthwhile emphasizing that the perturbative contribution to $\Pi_1^\mathcal{O}(0)$ has to be subtracted as it has already been included through the perturbative quark propagator in the presence of the external field (see Appendix).

Obviously, it is difficult to evaluate $\Pi_1^\mathcal{O}(0)$ directly from QCD at this stage. The QCD sum-rule method has been invoked to estimate $\Pi_1^\mathcal{O}(0)$ previously. The results given in Ref. [8] are $\chi a \approx -0.15 \text{GeV}^2$ and $\kappa a = \xi a \approx 0.1 \text{GeV}^4$. Recently, the authors of Ref. [21] have argued that the estimate of the vacuum tensor susceptibility χ in Ref. [8] is very rough, and their detailed analyses lead to a value for χ which has opposite sign and much larger (~ 4 times larger) magnitude. Given this uncertainty, we will consider the χ values in the following range

$$-0.5 \text{GeV}^2 \leq \chi a \leq 1.5 \text{GeV}^2 . \quad (32)$$

Since κ and ξ are small and the sum rules are relatively insensitive to their values, we use the values of κ and ξ given in Ref. [8] for simplicity.

The extraction of the nucleon tensor charge from the sum rules also requires the knowledge of the nucleon mass M_N , the couplings λ_1 and λ_2 , and the continuum threshold s_0 (for the sum rules in the usual formalism). These parameters can be determined from the corresponding mass sum rules. Here we take the central values obtained in Ref. [13]

$$M_N = 0.96 \text{GeV} , \quad \tilde{\lambda}_1 \tilde{\lambda}_2 = 0.41 \text{GeV}^6 , \quad s_0 = 1.69 \text{GeV}^2 , \quad (33)$$

for the sum rules from the mixed correlator and

$$M_N = 1.17 \text{GeV} , \quad \tilde{\lambda}_1^2 = 0.20 \text{GeV}^6 , \quad s_0 = 2.34 \text{GeV}^2 , \quad (34)$$

for the sum rules from the $1/2$ correlator. Note that the nucleon mass in Eq. (33) is very close to its experimental value. It is, however, much larger than the experimental value in Eq. (34). In the literature it is often found that the nucleon mass is fixed at its experimental value and the mass sum rules are used to determine the couplings and continuum threshold. This may introduce artificial errors. Moreover, when the same criteria are applied to the analyses of the nucleon mass sum rules, we find that the continuum threshold cannot be searched successfully when the nucleon mass is held fixed at its experimental value.

B. Results

Let us start from the sum rules in the usual formalism, Eqs.(19–20) and (26). The nucleon tensor charge g_T and the transition strength A_i are extracted from the sum rules, with the continuum threshold fixed at its value for the corresponding nucleon mass sum rule. We find that for the χ values interested here there does not exist valid Borel window for the sum rule Eq. (20) according to the criteria set above; it is thus invalid.




FIG. 1. Sum-rule predictions for the isovector nucleon tensor charge g_T^v as functions of χ . The solid and dashed curves correspond to the sum rules Eqs. (19) and (26), respectively. The dotted curve is also from the sum rule Eq. (26) but with $M_N = 0.94\text{GeV}$, $\tilde{\lambda}_1^2 = 0.26\text{GeV}^6$, and $s_0 = 2.3\text{GeV}^2$ as given in Ref. [8].

The sum rule Eq. (19) works only for a limited range of χ values. The predicted isovector (g_T^v) and isoscalar (g_T^s) nucleon tensor charges as functions of χ are plotted in Figs. 1 and 2, respectively. Both g_T^v and g_T^s increase as χ increases with essentially the same rate. This indicates that δu strongly depends on χ while δd has much weaker dependence on χ . One can see that $g_T^s > g_T^v > 0$, implying $\delta u > \delta d > 0$. We also find that the size of the valid Borel window shrinks with decreasing χ .

The sum rule Eq. (26) is valid for all the χ values considered here. Its predictions for the nucleon tensor charges are also given in Figs. 1 and 2. The predicted tensor charges increase as χ increases with a similar rate as those obtained from Eq. (19). The predictions again show $\delta u > \delta d > 0$, but δd has stronger χ dependence than that found from Eq. (19). For a given χ value, the size of the valid Borel window is in general larger than that for Eq. (19).

One notices that the predictions from Eq. (26) are significantly larger than those from Eq. (19). This partially attributes to the large nucleon mass and small coupling [see Eq. (34)] used in Eq. (26). To further illustrate this point, we also displayed in Figs. 1 and 2 the predictions obtained from the sum rule Eq. (26), with $M_N = 0.94\text{GeV}$, $\tilde{\lambda}_1^2 = 0.26\text{GeV}^6$, and $s_0 = 2.3\text{GeV}^2$ as quoted in Ref. [8]. In this case, the sum rule Eq. (26) only works for a limited range of χ values and the results are much closer to those from Eq. (19).

We now turn to the sum rules in the alternative formalism, Eqs. (22–23) and (27). As emphasized earlier, the continuum threshold s_0 must be treated as an unknown parameter to be extracted from the sum rules along with the tensor charges. We find that the sum




FIG. 2. Sum-rule predictions for the isoscalar nucleon tensor charge g_T^s as functions of χ . The solid and dashed curves correspond to the sum rules Eqs. (19) and (26), respectively. The dotted curve is also from the sum rule Eq. (26) but with $M_N = 0.94\text{GeV}$, $\tilde{\lambda}_1^2 = 0.26\text{GeV}^6$, and $s_0 = 2.3\text{GeV}^2$ as given in Ref. [8].

rule Eq. (23) fails to have a valid Borel window and is thus invalid. The sum rules Eqs. (22) and (27) are also invalid for the case of isoscalar tensor charge. For isovector tensor charge, the sum rule Eq. (19) only works for small χ values ($\chi \leq 0.1$) and its prediction is negative ($\sim -1.02 - -0.16$). These results are much smaller than the corresponding ones in the usual formalism. On the other hand, the sum rule Eq. (27) is only valid for large χ value ($\chi \geq 0.8$) and the predictions are slightly larger than those found from Eq. (26).

C. Discussion

To see how well the valid sum rules work, we have plotted in Fig. 3 the left- and right-hand sides of the sum rules Eqs. (19) and (26) for $\chi = 1.0$, with g_T^v and A_i extracted from the sum rules. The corresponding valid Borel windows and the relative contributions of the transition plus continuum and the highest order term in the OPE are displayed in Fig. 4. One notices that the sum rule Eq. (26) is valid in a broad Borel regime where the transition plus continuum contributes about 20% at the lower bound and the required 50% at the upper bound. The sum rule Eq. (19) has a much smaller valid Borel regime where the contributions from the transition and continuum are greater than 40% throughout the valid Borel window. On the other hand, the overlap of the two sides of Eq. (19) is better than that of Eq. (26). This pattern is seen at other χ values. It is therefore hard to compare the reliabilities of these two sum rules directly.




FIG. 3. The left- and right-hand sides of the sum rules Eqs. (19) and (26) as functions of the Borel mass for $\chi = 1.0$, with g_T^v and A_i extracted from the sum rules. The four curves correspond to the left- (solid curve) and right-hand (dashed curve) sides of Eq. (19) and the left- (dotted curve) and right-hand (dot-dashed curve) sides of Eq. (26), respectively

As shown in Eq. (15), the contribution from the transitions is in general a complicated function of the Borel mass, which is crudely modeled by a constant (A_i) in the usual formalism. This will change the curvature of the phenomenological side as a function of the Borel mass and may lead to errors to the extracted nucleon tensor charges. It is unclear how to reliably estimate the size of these errors within the criteria discussed in the present paper, though some estimates have been given in the study of other nucleon properties [16,22].

Moreover, the spectral parameters (mass, coupling, and continuum threshold) appearing in the nucleon mass sum rules also enter the sum rules for the nucleon tensor charge. This means that the uncertainties associated with these parameters will give rise to additional uncertainties, in addition to the uncertainties in the sum rules for the tensor charge themselves. This is a general drawback of the external field sum-rule approach, and we have demonstrated this point explicitly here. Clearly, one needs accurate knowledge of the vacuum spectral parameters in order to extract the tensor charges cleanly, which is difficult to achieve given the current implementation of the QCD sum-rule approach. Therefore, we expect large uncertainties associated with the predictions of the valid sum rules in the usual formalism presented above.

In the alternative formalism, the contribution of the transitions is absorbed into the continuum. It has been argued recently by one of us [16] that this alternative formalism may have the potential to improve the reliability of external-field sum-rule calculations, as the transitions are explicitly modeled by the modified continuum model. However, we have

found here that the sum rules in the alternative formalism fail in most cases interested here. This may indicate that the information contained in the truncated OPE is not enough and/or the transition and continuum give rise to too much background noise to isolate the information about the nucleon tensor charge. This might also suggest that the reason for the sum rules in the usual formalism to work is because the transition contribution is approximated by a constant.

We have emphasized throughout this paper that in the alternative formalism the continuum threshold has to be treated as a unknown parameter because the continuum model is modified relative to that in the usual formalism and contains the transition contribution. Fixing the continuum threshold at its value for the corresponding mass sum rule may introduce arbitrary artificial effect to the extracted tensor charges. In Ref. [8], the sum rule Eq. (27) is used in extracting the nucleon tensor charges but with the continuum threshold fixed at its value for the nucleon mass sum rule. However, we have found that this sum rule is invalid for $\chi a = -0.15\text{GeV}^2$ as used in Ref. [8], and hence the results obtained from this sum rule are unreliable.

In their analyses, the authors of Ref. [8] express the tensor charge as a function of the Borel mass by dividing both sides of Eq. (27) by $M_N \tilde{\lambda}_1^2 e^{-M_N^2/M^2}$ and find the tensor charge changes quickly with the Borel mass. This is actually an indication of the poor overlap of the two sides of the sum rule. To illustrate this point, we have plotted in Fig. 5 the left- and right-hand sides of Eq. (27), with the use of $M_N = 0.94\text{GeV}$, $\tilde{\lambda}_1^2 = 0.26\text{GeV}^6$, $\chi a = -0.15\text{GeV}^2$, and the continuum threshold fixed at $s_0 = 2.3\text{GeV}^2$ as quoted in Ref. [8].




FIG. 4. Relative contributions of the transition plus continuum and the highest order OPE terms to the sum rules as functions of the Borel mass. The dashed and dot-dashed curves correspond to the sum rules Eqs. (19) and (26), respectively.




FIG. 5. The left- (solid curve) and right-hand (dashed curve) sides of the sum rule (27) as functions of the Borel mass, with the use of $M_N = 0.94\text{GeV}$, $\tilde{\lambda}_1^2 = 0.26\text{GeV}^6$, $\chi a = -0.15\text{GeV}^2$, and the continuum threshold fixed at $s_0 = 2.3\text{GeV}^2$ as quoted in Ref. [8]. The sum rule is not optimized and $g_T^s = 0.3$ is adopted.

Here following Ref. [8], the sum rule is not optimized and the prediction for the nucleon tensor charge $g_T^s \simeq 0.3$ is obtained by simply taking $M^2 = 1\text{GeV}^2$. We see that the two sides of the sum rule does not match at all and they only coincide at the point $M^2 = 1\text{GeV}^2$, which is picked by hand. So, the prediction can be arbitrary as one may choose an arbitrary value for the Borel mass.

IV. SUMMARY AND CONCLUSION

In this paper, we have derived two new QCD sum rules for the nucleon tensor charge from the mixed correlator of spin-1/2 and spin-3/2 interpolating fields. Both the usual formalism and the alternative formalism were adopted to handle the transitions between the ground state nucleon and excited states. We analyzed these new sum rules, along with the chiral-odd sum rule obtained from the usual 1/2 correlator, with respect to specific criteria for monitoring the contaminations arising from the transitions and continuum and the convergence of the OPE. An emphasis was put on revealing the validity and reliability of various sum rules.

We found that in the usual formalism one of the two new sum rules is invalid and the other works only for certain values of χ . The predictions of the latter for moderate χ values ($0.5\text{ GeV}^2 < \chi a < 1.0\text{ GeV}^2$) are

$$\delta u \sim 0.67 - 1.06 , \quad \delta d \sim 0.23 - 0.24 . \quad (35)$$

This shows that δu has strong dependence on the vacuum tensor susceptibility χ and δd is insensitive to χ . The sum rule from the $1/2$ correlator is valid for all the χ values considered, and the predictions for the moderate χ are

$$\delta u \sim 3.48 - 3.89 , \quad \delta d \sim 0.12 - 0.18 . \quad (36)$$

It should be warned that the uncertainties quoted here only reflect the uncertainties from the χ value. There are large uncertainties associated with the usual formalism arising from the treatment of the transitions, the spectral parameters in the mass sum rules, and the vacuum condensates and other susceptibilities. Without evaluating these uncertainties, we cannot make a critical comparison with the results from other models and approaches as these uncertainties may change these results significantly. Therefore, the above results should be treated only as benchmark (at most). Nevertheless, our results appear to support $\delta u > \delta d > 0$. This behavior is different from that found in Rebs. [9–11] and the behavior of the quark spin structure of the nucleon ($\Delta u > 0$, $\Delta d < 0$).

The sum rules in the alternative formalism are invalid in most cases of interest. In particular, for isoscalar tensor charge all three sum rules are invalid. For isovector tensor charge, one of the two new sum rules is invalid for all the χ values, and the other only works marginally for small χ values and its predictions for the tensor charge are much smaller than those given in Eq. (35). The sum rule from the $1/2$ correlator works only for large χ values and its predictions are similar to those listed in Eq. (36). It is also found that the sum rule Eq. (27) with $\chi a = -0.15\text{GeV}^2$, studied in the previous works, is invalid.

In conclusion, the QCD sum rules for the nucleon tensor charge work only in limited cases and their predictions are expected to have large uncertainties. It is important to examine the validity and reliability of the sum rules before the extraction can be made. Without such an examination, the extracted results are likely to be arbitrary and misleading. To improve the performances of the sum rules for the nucleon tensor charge, one should have more accurate evaluation of the vacuum susceptibilities and vacuum condensates, more precise knowledge of the mass sum-rule spectral parameters, higher order terms in the OPE, and better approaches for treating the transitions.

ACKNOWLEDGMENTS

We thank Bob Jaffe for helpful discussions, and B. L. Ioffe, Kai Hennen and Hanxin He for useful correspondences. This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement #DF-FC02-94ER40818.

APPENDIX:

In this appendix, we give some ingredients for calculating the QCD side as obtained from carrying out the OPE. Further details of the technique can be found in the literature. As

usual, we work to the leading order in perturbative theory and neglect the up and down current quark masses, the isospin breaking effect, and the strangeness contribution. It is convenient to calculate the Wilson coefficients of the OPE in coordinate space and then perform the Fourier transformation to momentum space. To first order in the external field, the quark propagator without background gluon field can be expressed as

$$\begin{aligned}
\langle T[q_i^a(x)q_j^b(0)] \rangle_z &= \frac{i}{2\pi^2} \frac{\hat{x}_{ij}}{x^4} \delta^{ab} - \frac{\langle \bar{q}q \rangle}{12} \delta_{ij} \delta^{ab} + \frac{x^2}{192} \langle \bar{q}g_s \sigma \cdot Gq \rangle \delta_{ij} \delta^{ab} \\
&- \frac{x^4}{27648} \langle \bar{q}q \rangle \langle g_s^2 G^2 \rangle \delta_{ij} \delta^{ab} - \frac{g_q}{4\pi^2} Z^{\alpha\beta} \frac{[\hat{x}\sigma_{\alpha\beta}\hat{x}]_{ij}}{x^4} \delta^{ab} \\
&- \frac{g_q}{24} \chi \langle \bar{q}q \rangle Z^{\alpha\beta} (\sigma_{\alpha\beta})_{ij} \delta^{ab} - i \frac{g_q}{48} \langle \bar{q}q \rangle Z^{\alpha\beta} (\hat{x}\sigma_{\alpha\beta} + \sigma_{\alpha\beta}\hat{x})_{ij} \delta^{ab} \\
&+ \frac{g_q}{288} \langle \bar{q}q \rangle Z^{\alpha\beta} \left[x^2 (\kappa + 2\xi) \sigma_{\alpha\beta} + 2x_\beta x^\gamma (\kappa - \xi) \sigma_{\alpha\gamma} \right]_{ij} \delta^{ab} + \dots, \tag{A1}
\end{aligned}$$

where $\{i, j\}$ and $\{a, b\}$ are Dirac and color indices, respectively, and the fifth term is the perturbative propagator in the presence of the external field. Note that the sign in front of ξ differs from that given in Ref. [8]. Here we adopted the conventions of Itzykson and Zuber for $\epsilon_{\alpha\beta\rho\lambda}$ while Ref. [8] has not given the conventions explicitly.

In the fixed-point gauge, the quark propagator in the presence of background gluon field is given by

$$\begin{aligned}
\langle T[q_i^a(x)\bar{q}_j^b(0)] \rangle_z &= \frac{i}{32\pi^2} g_s G_{\rho\lambda}^n \frac{[\hat{x}\sigma^{\rho\lambda} + \sigma^{\rho\lambda}\hat{x}]_{ij}}{x^2} \left(\frac{\lambda^n}{2} \right)^{ab} \\
&+ \frac{1}{288} \langle \bar{q}q \rangle g_s G_{\rho\lambda}^n \left[x^2 \sigma^{\rho\lambda} - 2x_\delta x^\lambda \sigma^{\delta\rho} \right]_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} \\
&+ \frac{g_q}{64\pi^2} g_s G_{\rho\lambda}^n Z_{\alpha\beta} \left\{ -\frac{\sigma^{\rho\lambda} \hat{x} \sigma^{\alpha\beta} \hat{x} + \hat{x} \sigma^{\alpha\beta} \hat{x} \sigma^{\rho\lambda}}{x^2} \right. \\
&+ i \left(\gamma^\rho \sigma^{\alpha\beta} \gamma^\lambda - \gamma^\lambda \sigma^{\alpha\beta} \gamma^\rho \right) \left[\ln \left(-\frac{x^2 \Lambda^2}{4} \right) + 2\gamma_{\text{EM}} \right] \Big\}_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} \\
&- \frac{g_q}{6912} \langle g_s^2 G^2 \rangle Z_{\alpha\beta} \left[\ln \left(-\frac{x^2 \Lambda^2}{4} \right) + 2\gamma_{\text{EM}} \right] [\hat{x}\sigma^{\alpha\beta}\hat{x}]_{ij} \delta^{ab} \\
&+ i \frac{g_q}{288} \langle \bar{q}q \rangle g_s G_{\rho\lambda}^n \left[Z^{\rho\lambda} x^2 \hat{x} + 2Z^{\rho\delta} x^2 x^\lambda \gamma_\delta + 2i \tilde{Z}^{\delta\rho} x^2 x_\delta \gamma_5 \gamma^\lambda \right]_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} \\
&+ i \frac{g_q}{24} \chi \langle \bar{q}q \rangle g_s G_{\rho\lambda}^n Z_{\alpha\beta} x^\beta x^\lambda [\sigma^{\alpha\rho}]_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} \\
&+ \frac{g_q}{144} \chi \langle \bar{q}q \rangle g_s G_{\rho\lambda}^n \left(Z^{\rho\lambda} x^2 - Z^{\rho\delta} x_\delta x^\lambda \right) \delta_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} + \dots, \tag{A2}
\end{aligned}$$

where the third and fourth terms arise from the presence of both the external and background gluon fields. The last three terms have been ignored in Ref. [8], which gives rise to the differences in the last four terms in the sum rule Eq. (27). In the calculations of the terms proportional to ab , we have neglected the fourth and fifth terms of Eq. (A2). We have found that the inclusion of such terms only make tiny refinement to the sum rules.

We also need the following expression for the expectation value of the quark and gluon field product in the presence of the external field [15]

$$\begin{aligned} \langle q_i^a \bar{q}_j^b g_s G_{\alpha\beta}^n \rangle_z &= -\frac{1}{192} \langle \bar{q} g_s \sigma \cdot G q \rangle [\sigma_{\alpha\beta}]_{ij} \left(\frac{\lambda^n}{2} \right)^{ab} \\ &\quad - \frac{1}{16} \left[\kappa Z_{\alpha\beta} + \frac{1}{2} i \xi \epsilon_{\alpha\beta\rho\lambda} Z^{\rho\lambda} \gamma_5 \right]_{ij} \left(\frac{\lambda^n}{2} \right)^{ab}. \end{aligned} \quad (A3)$$

Now one can easily carry out the calculations of the correlators by expressing the correlators as [13]

$$\begin{aligned} \langle T[\eta_\mu^{1/2}(x) \bar{\eta}_\nu^{3/2}(0)] \rangle_z &= \\ &\epsilon^{abc} \epsilon^{a'b'c'} \left\{ \beta \gamma_\mu S_u^{aa'} \gamma_\nu \sigma_{\rho\lambda} \text{Tr} \left[S_d^{bb'} \sigma^{\rho\lambda} C S_u^{cc'T} C \right] + \gamma_\mu \gamma_5 S_u^{aa'} \gamma_\nu \sigma_{\rho\lambda} \text{Tr} \left[\gamma_5 S_d^{bb'} \sigma^{\rho\lambda} C S_u^{cc'T} C \right] \right. \\ &\quad - 2 \gamma_\mu \gamma_5 S_u^{aa'} \sigma_{\rho\lambda} C S_u^{cc'T} C \gamma_5 S_d^{bb'} \gamma_\nu \sigma^{\rho\lambda} - \gamma_\mu \gamma_5 S_u^{aa'} \sigma_{\rho\lambda} C S_d^{cc'T} C \gamma_5 S_u^{bb'} \gamma_\nu \sigma^{\rho\lambda} \\ &\quad \left. - 2 \beta \gamma_\mu S_u^{aa'} \sigma_{\rho\lambda} C S_u^{cc'T} C S_d^{bb'} \gamma_\nu \sigma^{\rho\lambda} - \beta \gamma_\mu S_u^{aa'} \sigma_{\rho\lambda} C S_d^{cc'T} C S_u^{bb'} \gamma_\nu \sigma^{\rho\lambda} \right\}, \end{aligned} \quad (A4)$$

$$\begin{aligned} \langle T[\eta^{1/2}(x) \bar{\eta}^{1/2}(0)] \rangle_z &= \\ &- \epsilon^{abc} \epsilon^{a'b'c'} \left\{ \beta^2 \gamma_5 S_u^{aa'} \gamma_5 \text{Tr} \left[C S_d^{cc'T} C S_u^{bb'} \right] + S_u^{aa'} \text{Tr} \left[C S_d^{cc'T} C \gamma_5 S_u^{bb'} \gamma_5 \right] \right. \\ &\quad + \beta \gamma_5 S_u^{aa'} \text{Tr} \left[C S_u^{cc'T} C S_d^{bb'} \gamma_5 \right] + \beta S_u^{aa'} \gamma_5 \text{Tr} \left[C S_u^{cc'T} C \gamma_5 S_d^{bb'} \right] \\ &\quad + S_u^{aa'} \gamma_5 C S_d^{cc'T} C \gamma_5 S_u^{bb'} + \beta \gamma_5 S_u^{aa'} \gamma_5 C S_d^{cc'T} C S_u^{bb'} \\ &\quad \left. + \beta S_u^{aa'} C S_d^{cc'T} C \gamma_5 S_u^{bb'} \gamma_5 + \beta^2 \gamma_5 S_u^{aa'} C S_d^{cc'T} C S_u^{bb'} \gamma_5 \right\}, \end{aligned} \quad (A5)$$

with the quark propagator $S_q^{ab}(x, 0; Z) \equiv \langle T[q^a(x) \bar{q}^b(0)] \rangle_z$ given above. Since only linear response of the correlators is concerned, one only needs to keep the terms linear in the external field.

Finally, we observe that the Borel transformation of $\ln(-q^2)/q^2$ has not been carried out correctly in Ref. [8]. So, we give the correct formula [19]

$$\mathcal{B} \left[\frac{\ln(-q^2)}{q^2} \right] = -\ln(M^2) + \gamma_{\text{EM}}, \quad (A6)$$

from which we get

$$\mathcal{B} \left[\left(M_N^2 - q^2 \right) \frac{\ln(\Lambda^2 / -q^2)}{q^2} \right] = M_N^2 \left[\ln \left(\frac{M^2}{\Lambda^2} \right) - \gamma_{\text{EM}} \right] - M^2. \quad (A7)$$

It is then easy to show that the second term of Eq. (20) in Ref. [8] is not the correct Borel transform of the second term of Eq. (17).

REFERENCES

- [1] R. L. Jaffe and X. Ji, Phys. Rev. Lett, **71**, 2547 (1993).
- [2] R. L. Jaffe, “*Spin, Twist and Hadron Structure in Deep Inelastic Processes*”, Lectures presented at the International School of Nucleon Structure, Erice, Italy August 3-10, 1995, Report No. MIT-CTP-2506, hep-ph/9602236, and references therein.
- [3] J. Ralston and D. E. Soper, Nucl. Phys. **B152**, 109 (1979).
- [4] RSC Collaboration, G. Bunce *et al.*, Particle World **3**, 1 (1992); Proposal on Spin Physics using RHIC Polarized Collider, R5 (August 1992), approved October 1993.
- [5] The HERMES proposal to HERA (1993).
- [6] COMPASS Collaboration (G. Baum *et al.*), Report No. CERN-SPSLC-96-14 (1996).
- [7] H. X. He and X. Ji, Phys. Rev. D **52**, 2960 (1995).
- [8] H. X. He and X. Ji, Phys. Rev. D **54**, 6897 (1996).
- [9] H.-C. Kim, M. V. Polyakov, and K. Goeke, Phys. Rev. D **53**, 4715 (1996); Phys. Lett. **B387**, 577 (1996).
- [10] I. Schmidt and J. Soffer, Report No. hep-ph/9703411.
- [11] S. Aoki, M. Doui, T. Hatsuda, and Y. Kuramashi, Report No. hep-lat/9608115.
- [12] B. L. Ioffe, Nucl. Phys. **B188**, 317 (1981).
- [13] D. B. Leinweber, Ann. Phys. (N.Y.) **254**, 328 (1997).
- [14] F. X. Lee, D. B. Leinweber, and X. Jin, Phys. Rev. D **55**, 4066 (1997).
- [15] B. L. Ioffe and A. V. Smilga, Nucl. Phys. **B232**, 109 (1984).
- [16] X. Jin, Phys. Rev. D **55**, 1693 (1997).
- [17] Y. Chung, H. G. Dosch, M. Kremer, and D. Schall, Nucl. Phys. **B197**, 55 (1982).
- [18] X. Jin and J. Tang, Report No. MIT-CTP-2601, hep-ph/9701230, Phys. Rev. D **56**, (1997) in press.
- [19] S. L. Wilson, J. Pasupathy, and C. B. Chiu, Phys. Rev. D **36**, 1451 (1987).
- [20] X. Jin and D. B. Leinweber, Phys. Rev. C **52**, 3344 (1995); M. J. Iqbal, X. Jin, and D. B. Leinweber, Phys. Lett. **B386**, 55 (1996); R. J. Furnstahl, X. Jin, and D. B. Leinweber, Phys. Lett. **B387**, 253 (1996).
- [21] V. M. Belyaev and A. Oganesian, Phys. Lett. **B395**, 307 (1997).
- [22] B. L. Ioffe, Phys. At. Nucl. **58**, 1408 (1995).